

Human Reproduction, Vol 8, 393-395, (1993) by Oxford University Press

IMPROVEMENT OF CLINICAL PREGNANCY RATE AND IMPLANTATION RATE OF IN- VITRO FERTILIZATION-EMBRYO TRANSFER PATIENTS BY USING METHYLPREDNISONE

E Polak de Fried, L Blanco, S Lancuba and RH Asch
Centro de Salud Reproductiva (CER), Buenos Aires, Argentina.

A prospective non-randomized study was undertaken to test whether immunosuppression improves implantation and pregnancy rates in an in- vitro fertilization-embryo transfer (IVF-ET) programme in patients with tubal factor infertility. Treatment involved ovarian stimulation, transvaginal oocyte retrieval, IVF-ET, and assessment of short-term administration of large doses of corticosteroids (60 mg of methyl- prednisone x 4 days). When compared to the group that did not receive immunosuppressive doses of methylprednisone (group A; mean age 31.85 +/- 4.09 years), those subjects who were treated (group A2) showed a statistically significant increase in pregnancy ($P < 0.01$) and take home baby rate ($P < 0.01$). Similar results were observed in subjects who received corticosteroids in their first IVF-ET attempt (group B; mean age 34.32 +/- 4.98 years). Our results suggest that immunosuppressive doses of corticosteroids administered for a short period of time to patients undergoing IVF-ET could significantly improve the implantation and pregnancy rates. Possible mechanisms of action of corticosteroids are proposed.